

1-2016

## Geographically Isolated Wetlands are Part of the Hydrological Landscape

Mark Rains

*University of South Florida, [mrains@usf.edu](mailto:mrains@usf.edu)*

S. G. Leibowitz

*US EPA National Health and Environmental Effects Research Laboratory*

M. J. Cohen

*University of Florida*

I. F. Creed

*Western University*

H. E. Golden

*US EPA National Exposure Research Laboratory*

*See next page for additional authors*

Follow this and additional works at: [https://digitalcommons.usf.edu/geo\\_facpub](https://digitalcommons.usf.edu/geo_facpub)



Part of the [Geology Commons](#)

---

### Scholar Commons Citation

Rains, Mark; Leibowitz, S. G.; Cohen, M. J.; Creed, I. F.; Golden, H. E.; Jawitz, J. W.; Kalla, P.; Lane, C. R.; Lang, M. W.; and McLaughlin, D. L., "Geographically Isolated Wetlands are Part of the Hydrological Landscape" (2016). *School of Geosciences Faculty and Staff Publications*. 1093.  
[https://digitalcommons.usf.edu/geo\\_facpub/1093](https://digitalcommons.usf.edu/geo_facpub/1093)

This Response or Comment is brought to you for free and open access by the School of Geosciences at Digital Commons @ University of South Florida. It has been accepted for inclusion in School of Geosciences Faculty and Staff Publications by an authorized administrator of Digital Commons @ University of South Florida. For more information, please contact [digitalcommons@usf.edu](mailto:digitalcommons@usf.edu).

---

**Authors**

Mark Rains, S. G. Leibowitz, M. J. Cohen, I. F. Creed, H. E. Golden, J. W. Jawitz, P. Kalla, C. R. Lane, M. W. Lang, and D. L. McLaughlin

# Geographically isolated wetlands are part of the hydrological landscape

M. C. Rains,<sup>1\*</sup>  
S. G. Leibowitz,<sup>2</sup>  
M. J. Cohen,<sup>3</sup> I. F. Creed,<sup>4</sup>  
H. E. Golden,<sup>5</sup> J. W. Jawitz,<sup>6</sup>  
P. Kalla,<sup>7</sup> C. R. Lane,<sup>5</sup>  
M. W. Lang<sup>8</sup> and  
D. L. McLaughlin<sup>9</sup>

<sup>1</sup> School of Geosciences, University of South Florida, Tampa, FL, 33620, USA

<sup>2</sup> US EPA National Health and Environmental Effects Research Laboratory, Corvallis, OR, 97333, USA

<sup>3</sup> School of Forest Resources and Conservation, University of Florida, Gainesville, FL, 32611, USA

<sup>4</sup> Department of Biology, Western University, London, ON, N6A 5B7, Canada

<sup>5</sup> US EPA National Exposure Research Laboratory, Cincinnati, OH, 45268, USA

<sup>6</sup> Soil and Water Science Department, University of Florida, Gainesville, FL 32611, USA

<sup>7</sup> US EPA Region 4, Athens, GA, 30606, USA

<sup>8</sup> Department of Geographical Sciences, University of Maryland, College Park, MD, 20742, USA

<sup>9</sup> Department of Forest Resources and Environmental Conservation, Virginia Tech, Blacksburg, VA, 24061, USA

\*Correspondence to:

MC Rains, School of Geosciences, University of South Florida, Tampa, FL 33620, USA.

E-mail: mrains@usf.edu

The copyright line for this article was changed on 10 February 2016 after original online publication.

Received 26 June 2015

Accepted 3 July 2015

## Introduction

Since the US Supreme Court's 2001 *SWANCC* case (531 US 159), there has been significant focus on whether Clean Water Act (CWA) protections should be extended to so-called geographically isolated wetlands (GIWs); wetlands that are surrounded by uplands and lack readily apparent surface water connections to downgradient waters (Downing *et al.*, 2003; Leibowitz and Nadeau, 2003; Tiner, 2003a, b; see Mushet *et al.* (2015) for a history and critique of this term). Following the US Supreme Court's 2006 *Rapanos* case (547 US 715), interest in GIWs increased, with a more recent emphasis on the roles surface and subsurface hydrological flows might play in connecting GIWs to downgradient waters at the landscape scale (Downing *et al.*, 2007; Nadeau and Rains, 2007; Leibowitz *et al.*, 2008). One key outcome from *Rapanos* comes from the opinion penned by Justice Anthony Kennedy, which states that non-adjacent wetlands, including non-adjacent GIWs, can be waters of the USA (WOUS) subject to regulation under the CWA if they, either individually or cumulatively, have a 'significant nexus' with the chemical, physical, and/or biological integrity of other, more traditionally defined WOUS (e.g. navigable waters). In other words, a GIW is a WOUS if it is connected to a downgradient WOUS, and this connection substantively contributes to the chemical, physical, and/or biological integrity of that downgradient WOUS.

The US Environmental Protection Agency recently completed a review of peer-reviewed literature, seeking to synthesize existing scientific understanding of how wetlands and streams, individually or in aggregate, affect the chemical, physical, and biological integrities of downstream waters (US Environmental Protection Agency, 2015). The report concludes that all wetlands located on floodplains and/or within riparian areas have significant chemical, physical, and/or biological connections with downgradient WOUS. The report is more equivocal about those other wetlands, including the vast majority of GIWs, which are not located on floodplains and/or riparian areas. Instead, the report concludes that these wetlands occur along a continuum of connectivity, with a great deal of spatial heterogeneity and temporal variability, and that a lack of knowledge makes any generalization difficult. The report cites more than 1200 peer-reviewed papers, suggesting that the problem is not a lack of general knowledge about wetlands and waters but, rather, a lack of specific knowledge on the roles that GIWs might play in controlling the chemical, physical, and/or biological integrities of downgradient WOUS. Therefore, the report concludes that additional research focused on the frequency, magnitude, timing, duration, and rate of fluxes from GIWs to downgradient waters is needed to improve the US Environmental

Protection Agency's abilities to 'identify waters of national importance and maintain the long-term sustainability and resiliency of valued water resources'. Towards these ends, a broad range of research is needed, from field and numerical modelling studies that evaluate connectivity and better elucidate functional relationships between GIWs and downgradient waters to the development of new conceptual frameworks that can be used to generate hypotheses regarding how these systems vary over space and time. The latter is the focus of this commentary, with a specific focus on the effects of GIWs on flows in downgradient waters.

### GIWs as Nodes in Hydrologic Networks

Hydrological flowpaths connect landscapes in four dimensions – longitudinal, lateral, vertical, and through time. This four-dimensional hydrological connectivity, operating at local to landscape scales, is a basic tenet of freshwater ecology (Ward, 1989). Hydrological flowpaths are extensive and dynamic, connecting landscapes within watersheds (McDonnell, 2013) and across watershed divides (see Sun *et al.*, 1997 and references therein). Fluxes of water along these hydrological flowpaths occur at varying frequencies, magnitudes, timings, durations, and rates, which are primarily determined by climate, geology, and topography (Winter, 2001; Wolock *et al.*, 2004; Devito *et al.*, 2005; Wigington *et al.*, 2013; Park *et al.*, 2014) and collectively control the physical integrity of downgradient waters (Nadeau and Rains, 2007). GIWs distributed throughout the landscape intercept and interact with water that flows along these flowpaths, and these GIWs are therefore integrally connected to uplands, other wetlands, and downgradient waters.

Conceptually, this hydrological landscape is a network, with GIWs as nodes – receiving, storing, and sending water – and flowpaths as edges – transmitting water (Figure 1). In the GIWs, flows are modulated by the performance of lag, sink, and source functions (Table I; Leibowitz *et al.*, 2008; US Environmental Protection Agency, 2015). Lag functions delay the flow of water to downgradient waters and include local surface water and groundwater storage (Haag *et al.*, 2005; Gleason *et al.*, 2007; Lane and D'Amico, 2010) and exchange (Min *et al.*, 2010; Nilsson *et al.*, 2013; McLaughlin *et al.*, 2014), with the latter regulating water tables and enhancing or reducing surface water and groundwater storage depending on the direction of the exchange. Sink functions reduce the flow of water to downgradient waters and include evapotranspiration (Sun *et al.*, 2002; Towler *et al.*, 2004; Hammersmark *et al.*, 2010) and deep groundwater recharge (Sinclair, 1977; Wood and Sanford, 1995; Rains, 2011). Lag and

sink functions can act in concert to more greatly affect wetland response to variable precipitation (Rosenberry and Winter, 1997). Collectively, these lag and sink functions modulate the source function that can contribute flow to downgradient waters by surface water and shallow groundwater outflow (Leibowitz and Vining, 2003; Rains *et al.*, 2006; Sass and Creed, 2008; Wilcox *et al.*, 2011; Golden *et al.*, 2015). Along the flowpaths, flows are further altered by interactions among flowpath length, gradient, resistance or conductance (e.g. hydraulic conductivity and surface roughness), and leakage (e.g. evapotranspiration and deep recharge). As water flows through this network, the frequency, magnitude, timing, duration, and rate of flows are all modulated by myriad interactions occurring in GIWs and along flowpaths.

### Network-Scale Effects of GIWs on Flow Generation

While an individual GIW can affect local-scale hydrology, its effect on landscape-scale hydrology is likely negligible. However, the cumulative effect of many GIWs can play an important role in landscape-scale hydrology by regulating the frequency, magnitude, timing, duration, and rate of flows to downgradient waters (Ogawa and Male, 1986; Hey and Philippi, 1995; Cohen and Brown, 2007; Golden *et al.*, 2015). This cumulative effect emerges from lag, sink, and source functions resulting in time-varying flows being directed towards downgradient waters along overland (Leibowitz and Vining, 2003; Wilcox *et al.*, 2011), shallow subsurface (Rains *et al.*, 2006; van der Kamp and Hayashi, 2009), and deep groundwater (Winter, 1999; Rains, 2011) flowpaths.

The GIWs and the flowpaths that connect them to downgradient waters exist along a hydrologically dynamic continuum (Euliss *et al.*, 2004; Cohen *et al.*, In Review). Geology and topography are spatially heterogeneous but temporally fixed. However, climate is variable, and its effects vary annually, seasonally, and episodically. Therefore, the degree to which lag, sink, and source functions are performed and the flowpaths along which water is directed to downgradient waters are strongly dependent on both current and antecedent conditions (Rains *et al.*, 2006; Pyzoha *et al.*, 2008). When lag and sink functions dominate and/or water is directed from GIWs to downgradient waters along shallow subsurface or deep groundwater flowpaths, then downgradient flows might be delayed or diminished. In contrast, when source functions dominate and/or water is directed from GIWs to downgradient waters along overland flowpaths, then downgradient flows might be hastened or enhanced. Therefore,




Figure 1. (a) A watershed with GIWs and other waterbodies conceptualized as (b) a network with GIWs and other waterbodies as nodes – receiving, storing, and sending water – and flowpaths as edges – transmitting water. GIW, geographically isolated wetland

Table I. Hydrologic functions of geographically isolated wetlands

Function	Type	Description
Storage	Lag	Storage of surface water and/or shallow groundwater. Partially controls other hydrologic functions. Especially pronounced in depressional GIWs.
Exchange	Lag	Exchange of surface water and groundwater, thereby regulating water table variation because of bidirectional exchanges (recharge and discharge) at local scales.
Evapotranspiration	Sink	Enhanced evapotranspiration because of prolonged presence of surface water and/or shallow groundwater. Potentially an important watershed-scale loss of water.
Deep recharge	Sink	Enhanced deep groundwater recharge because of local topographic lows that hold surface water and/or shallow groundwater. Potentially an important watershed-scale loss of water.
Flow generation	Source	Especially pronounced in depressional GIWs. Alteration of the frequency, magnitude, timing, duration, and rate of outflows to downgradient wetlands and waters because of the combined effects of the lag and sink functions.

GIW, geographically isolated wetland

antecedent conditions exert a substantial control on travel times, with water entering a GIW being directed along slow subsurface flowpaths, rapid surface flowpaths, to or from adjacent shallow groundwater, or to the atmosphere or deep groundwater storage depending upon those antecedent conditions.

The cumulative effect results from water flowing from many GIWs to downgradient waters along a continuum of travel lengths and times, varying by GIW and over

time. At a given moment in time, there might be no flow from some GIWs, relatively slow subsurface flow from other GIWs, and relatively rapid surface flow from still other GIWs. The cumulative effect of the many GIWs on downgradient streamflows emerges from the convolution of these travel times (Cohen *et al.*, In Review; Figure 2). In this convolution, time-varying flows – or the lack thereof – from each GIW cumulatively contribute to the maintenance of the natural flow regime (Poff *et al.*,




Figure 2. At a given moment in time, the effects of GIWs on downgradient hydrographs emerge from the convolution of the continuum of travel times between the portfolio of GIWs in the network and the downgradient water. The result is a component of the hydrograph composed of the time-varying contributions from each GIW in the network, which could collectively play important roles in maintaining the natural flow regime. Modified from Cohen *et al.* (In Review). GIW, geographically isolated wetland

1997). Because these flows are time varying, the effect on downgradient hydrographs is not fully realized until all GIWs have gone through complete annual and inter-annual cycles of connectivity (Phillips *et al.*, 2011), so altering any component of the convolved hydrological response could change the natural flow regime, with potential impacts to downgradient waters.

### Human Alterations to GIWs

Human alterations to GIWs can affect lag, sink, and source functions, thereby altering the convolved, watershed-scale hydrologic response. For example,

sediments can be deposited in GIWs – either by direct placement or indirect sediment-laden discharge from the contributing basin – reducing storage capacity, sometimes by as much as 100% (Luo *et al.*, 1997, 1999; Fenstermacher *et al.*, 2014). Direct drainage of GIWs can result in the disproportionate loss of small GIWs and those distant from the stream network at the landscape scale (Lang *et al.*, 2012), which can increase the runoff efficiency between the remaining wetlands and downgradient waters (Van Meter and Basu, 2015). Direct drainage also can alter the hydraulic gradients that arise between GIWs and adjacent uplands, thereby altering the surface water and groundwater exchange

between GIWs and adjacent uplands (McLaughlin and Cohen, 2013). Regional groundwater pumping can lower hydraulic heads, resulting in enhanced groundwater recharge from overlying GIWs (Haag *et al.*, 2005; Lee *et al.*, 2009; Haag and Pfeiffer, 2012).

Human alterations to flowpaths also can affect the convolved hydrologic response. Ditching and tiling (e.g. for agricultural purposes) increase drainage efficiency, with the result commonly being that water is quickly routed into and/or out of GIWs (Randall *et al.*, 1997; Min *et al.*, 2010; Boland-Brien *et al.*, 2014). This often more directly links otherwise remote GIWs to stream systems (Gamble *et al.*, 2007) and has a significant effect on downgradient streamflows (Cohen and Brown, 2007; Babbar-Sebens *et al.*, 2013). Even in the absence of direct ditching or tiling, changes in land use and/or land cover can alter flowpath dynamics, as the mechanical destruction of soil structure and the homogenization of microtopography in agricultural settings can increase runoff efficiency into and out of GIWs (Euliss and Mushet, 1996; van der Kamp *et al.*, 2003; Pyke and Marty, 2005; Tsai *et al.*, 2007; McDonough *et al.*, 2014), and preferential surface water flowpaths can be established through animal-mediated soil compaction along trails often terminating at GIWs (e.g. Tanner *et al.*, 1984; van der Kamp *et al.*, 2003; Franzluebbers *et al.*, 2012). In addition, increases in impervious surfaces in urban settings may also increase runoff efficiency, thereby contributing to downstream hydrograph ‘flashiness’ (Walsh *et al.*, 2012; Faulkner, 2004). Conversely, the restoration of GIWs may moderate flows (McDonough *et al.*, 2014).

### Implications for Future Research and Policy

The extensive and dynamic hydrological flowpaths connect landscapes in four dimensions is well known as regards to stream networks (Ward, 1989; Nadeau and Rains, 2007; McDonnell, 2013). More poorly known and understood is the role that GIW nodes and related flowpath edges play in the functioning of the broader hydrological network, including the stream network. This represents a critical knowledge gap, especially in archetypal GIW-dominated landscapes (e.g. vernal pools, prairie potholes, and Carolina bays) where the number of these GIW nodes and related flowpath edges is large (Semlitsch and Bodie, 1998).

Does recognition of GIWs as nodes within the hydrological network mean that there is a significant nexus between the nodes and the chemical, physical, and/or biological integrities of downgradient waters? Whether this constitutes a significant nexus is a policy decision that we do not purport to advance here; nevertheless, GIWs certainly perform lag, sink, and source functions that can influence the chemical,

physical, and/or biological integrities of downgradient waters, especially when considered in aggregate (Ogawa and Male, 1986; Hey and Philippi, 1995; Bullock and Acreman, 2003; Cohen and Brown, 2007; US Environmental Protection Agency, 2015). But few studies have sought to discern the specific effects of GIWs on downgradient waters (McLaughlin *et al.*, 2014; Golden *et al.*, 2015). Therefore, there remains a lack of general agreement on the roles that GIWs play in landscape-scale hydrology (US Environmental Protection Agency, 2015). Understanding the emergent properties of GIWs at the landscape-scale requires that we consider more than just the typical behaviour of a GIW or given class of GIW. Rather, it requires that we focus instead on the aggregate effects of a portfolio of functions and behaviours expressed by a network of GIWs and GIW complexes (Figure 2; Cohen *et al.*, In Review).

An initial step towards improving our understanding of the aggregate effects of GIWs is the development of a classification system that can be used to define regions or conditions under which GIWs have expected behaviours that can be studied in aggregate, much like the concept of hydrological landscapes (Winter, 2001; Wolock *et al.*, 2004; Wigington *et al.*, 2013), although defined at scales and including factors more appropriate for the study of GIWs (Rains *et al.*, 2008). The next step is to place an increasing emphasis on regional-scale data collection, including both field data and remote-sensing data. As regards to the latter, improvements in the sensitivity and temporal resolution of commonly available datasets might be necessary to map GIWs and related flowpaths, given that remote-sensing datasets traditionally used to map aquatic resources (e.g. aerial photographs) may have significant limitations when applied to GIWs (Lang *et al.*, 2012; Yang and Chu, 2012; Lang *et al.*, 2013). The final step is improving the sensitivity and accessibility of modelling and analytical tools that can be used to evaluate the aggregate effects of the portfolio of GIWs that emerge at the watershed scale. This might require the development of new model approaches with an explicit focus on the roles that GIWs play at the network scale (e.g. McLaughlin *et al.*, 2014). Alternatively, this might instead only require the adaptation of existing models to better describe the fine-scale surface–subsurface interactions that characterize connectivity between GIWs and the broader hydrological landscape (Golden *et al.*, 2014).

A concerted effort such as this could transform our understanding of watershed-scale hydrology, facilitating a better understanding of the roles played by GIWs and how these roles change depending upon spatial heterogeneity and temporal variability. A concerted effort such as this also would be timely given the ongoing debate about the geographic extent of the CWA, particularly as it relates to non-navigable, intrastate waters, including

GIWs. Under the current rule, non-adjacent wetlands, including non-adjacent GIWs, require case-by-case determinations of significant nexus if they are members of five subcategories (e.g. western vernal pools, prairie potholes, Delmarva Bays, Carolina Bays, or pocosins) or are within 1219 m of the ordinary high water mark or high tide line (US Army Corps of Engineers/US Environmental Protection Agency, 2015). A concerted effort such as this would improve upon the scientific understanding underlying the policy (e.g. US Environmental Protection Agency, 2015) and facilitate the development of improved policy and/or regulatory guidance (e.g. 80 FR 37054), better enabling decision-making regarding the geographic extent of WOUS subject to regulation under the CWA.

## Acknowledgements

The idea for this paper originated during a ‘Geographically Isolated Wetlands Research Workshop’ convened and co-hosted by the US Environmental Protection Agency Office of Research and Development and the Joseph W. Jones Ecological Research Center and held in Newton, GA, 18–21 November 2013 ([http://www.jonesctr.org/education\\_and\\_outreach/publications/GIW\\_Workshop\\_Summary\\_2013\\_v6.3.pdf](http://www.jonesctr.org/education_and_outreach/publications/GIW_Workshop_Summary_2013_v6.3.pdf)). We gratefully acknowledge all of the hosts and participants of that workshop. The information in this document has been funded in part by the US Environmental Protection Agency. This article has been subjected to US Environmental Protection Agency review and has been approved for publication. The views expressed in this article are those of the authors and do not necessarily reflect the views or policies of the US Environmental Protection Agency. Mention of trade names or commercial products does not constitute endorsement or recommendation for use.

## References

Babbar-Sebens M, Barr RC, Tedesco LP, Anderson M. 2013. *Spatial identification and optimization of upland wetlands in agricultural watersheds. Ecological Engineering* 52: 130–142.

Boland-Brien SJ, Basu NB, Schilling KR. 2014. Homogenization of spatial patterns of hydrologic response in artificially drained agricultural catchments. *Hydrological Processes* 28: 5010–5020.

Bullock A, Acreman M. 2003. The role of wetlands in the hydrological cycle. *Hydrology and Earth System Sciences* 7: 358–389.

Cohen MJ, Brown MT. 2007. A model examining hierarchical wetland networks for watershed stormwater management. *Ecological Modelling* 201: 179–193.

Cohen MJ, Creed IF, Alexander L, Basu N, Calhoun A, Craft C, D’Amico E, DeKeyser E, Fowler L, Golden HE, Jawitz JW, Kalla P, Kirkman LK, Lane CR, Lang M, Leibowitz SG, Lewis DB, Marton J, McLaughlin DL, Mushet D, Raanan-Kiperwas H, Rains MC, Smith

L, Walls S. In Review. Do geographically isolated wetlands impact landscape functions? *Proceedings of the National Academy of Sciences*.

Devito KJ, Creed IF, Gan T, Mendoza C, Petrone R, Silins U, Smerdon B. 2005. A framework for broad-scale classification of hydrologic response units on the Boreal Plain: is topography the first or last thing to consider? *Hydrological Processes* 19: 1705–1714.

Downing DM, Winer C, Wood LD. 2003. Navigating through Clean Water Act jurisdiction: a legal review. *Wetlands* 23: 475–493.

Downing D, Nadeau T-L, Kwok R. 2007. Technical and scientific challenges in implementing Rapanos’ ‘Water of the United States’. *Natural Resources and Environment* 22(42–45): 62–63.

Euliss NH Jr, Mushet DM. 1996. Water-level fluctuation in wetlands as a function of landscape condition in the prairie pothole region. *Wetlands* 16: 587–593.

Euliss N, LaBaugh J, Fredrickson L, Mushet D, Laubhan M, Swanson G, Winter T, Rosenberry D, Nelson R. 2004. The wetland continuum: a conceptual framework for interpreting biological studies. *Wetlands* 24: 448–458.

Faulkner S. 2004. Urbanization impacts on the structure and function of forested wetlands. *Urban Ecosystems* 7: 89–106.

Fenstermacher D, Rabenhorst M, Lang M, McCarty G, Needelman B. 2014. Distribution, morphometry, and land use of Delmarva Bays. *Wetlands* 34: 1219–1228.

Franzluebbers AJ, Stuedemann JA, Franklin DH. 2012. Water infiltration and surface-soil structural properties as influenced by animal traffic in the Southern Piedmont USA. *Renewable Agriculture and Food Systems* 27: 256–265.

Gamble D, Grody E, Undercoffer J, Mack JJ, Micacchion M. 2007. *An Ecological and Functional Assessment of Urban Wetlands in Central Ohio. Volume 2: Morphometric Surveys, Depth–Area–Volume Relationships and Flood Storage Function*, Ohio EPA Technical Report WET/2007-3B. Ohio Environmental Protection Agency, Wetland Ecology Group, Division of Surface Water: Columbus.

Gleason RA, Tangen BA, Laubhan MK, Kermes KE, Euliss NH Jr. 2007. Estimating water storage capacity of existing and potentially restorable wetland depressions in a subbasin of the Red River of the north. US Geological Survey Open-File Report 2007–1159.

Golden HE, Lane CR, Amatya DM, Bandilla KW, Raanan Kiperwas H, Knightes CD, Ssegane H. 2014. Hydrologic connectivity between geographically isolated wetlands and surface water systems: a review of select modeling methods. *Environmental Modelling and Software* 53: 190–206.

Golden HE, Sander SA, Lane CR, Zhao C, Price K, D’Amico E, Christensen JR. 2015. Relative effects of geographically isolated wetlands on streamflow: a watershed-scale analysis. *Ecohydrology*. DOI:10.1002/eco.1608.

Haag KH, Pfeiffer W. 2012. Flooded area and plant zonation in isolated wetlands in well fields in the northern Tampa Bay region, Florida, Following Reductions in Groundwater-Withdrawal Rates. US Geological Survey Scientific Investigations Report 2012–5039.

Haag KH, Lee TM, Herndon DC. 2005. Bathymetry and vegetation in isolated marsh and cypress wetlands in the northern Tampa Bay area, 2000–2004. US Geological Survey Scientific Investigations Report 2005–5109.

Hammersmark CT, Dobrowski S, Rains MC, Mount JF. 2010. Simulated effects of stream restoration on herbaceous vegetation distribution. *Restoration Ecology* 18: 882–893.

Hey DL, Philippi NS. 1995. Flood reduction through wetland restoration: the Upper Mississippi River Basin as a case history. *Restoration Ecology* 3: 4–17.

van der Kamp G, Hayashi M. 2009. Groundwater-wetland ecosystem interaction in the semiarid glaciated plains of North America. *Hydrogeology Journal* 17: 203–214.



- van der Kamp G, Hayashi M, Gallen D. 2003. Comparing the hydrology of grassed and cultivated catchments in the semi-arid Canadian prairies. *Hydrological Processes* 17: 559–575.
- Lane C, D'Amico E. 2010. Calculating the ecosystem service of water storage in isolated wetlands using LiDAR in north-central Florida, USA. *Wetlands* 30: 967–977.
- Lang M, McDonough O, McCarty G, Oesterling R, Wilen B. 2012. Enhanced detection of wetland–stream connectivity using LiDAR. *Wetlands* 32: 461–473.
- Lang M, McCarty G, Oesterling R, Yeo I-Y. 2013. Topographic metrics for improved mapping of forested wetlands. *Wetlands* 33: 141–155.
- Lee TM, Haag KH, Metz PA, Sacks LA. 2009. Comparative hydrology, water quality, and ecology of selected natural and augmented freshwater wetlands in West-Central Florida. US Geological Survey Professional Paper 1758.
- Leibowitz SG, Nadeau T-L. 2003. Isolated wetlands: state-of-the-science and future directions. *Wetlands* 23: 663–684.
- Leibowitz SG, Vining KC. 2003. Temporal connectivity in a prairie pothole complex. *Wetlands* 23: 13–25.
- Leibowitz SG, Wigington PJ Jr, Rains MC, Downing DM. 2008. Non-navigable streams and adjacent wetlands: addressing science needs following the Supreme Court's Rapanos decision. *Frontiers in Ecology and the Environment* 6: 364–371.
- Luo H-R, Smith LM, Allen BL, Haukos DA. 1997. Effects of sedimentation on playa wetland volume. *Ecological Applications* 7: 247–252.
- Luo H-R, Smith LM, Haukos DA, Allen BL. 1999. Sources of recently deposited sediments in playa wetlands. *Wetlands* 19: 176–181.
- McDonnell JJ. 2013. Are all runoff processes the same? *Hydrological Processes* 27: 4103–4111.
- McDonough OT, Lang MW, Hosen JD, Palmer MA. 2014. Surface hydrologic connectivity between Delmarva Bay wetlands and nearby streams along a gradient of agricultural alteration. *Wetlands* 35: 41–53.
- McLaughlin DL, Cohen MJ. 2013. Realizing ecosystem services: wetland hydrologic function along a gradient of ecosystem condition. *Ecological Applications* 23: 1619–1631.
- McLaughlin DL, Kaplan DA, Cohen MJ. 2014. A significant nexus: geographically isolated wetlands influence landscape hydrology. *Water Resources Research* 50: 7153–7166. DOI:10.1002/2013WR015002.
- Min JH, Perkins DB, Jawitz JW. 2010. Wetland–groundwater interactions in subtropical depression wetlands. *Wetlands* 30: 997–1006.
- Mushet DM, Calhoun AJK, Alexander LC, Cohen MJ, DeKeyser ES, Fowler L, Lane CR, Lang MW, Rains MC, Walls SC. 2015. Geographically isolated wetlands: rethinking a misnomer. *Wetlands*. DOI:10.1007/s13157-015-0631-9.
- Nadeau TL, Rains MC. 2007. Hydrological connectivity of headwaters to downstream waters: introduction to the featured collection. *Journal of the American Water Resources Association* 43: 1–4.
- Nilsson KA, Rains MC, Lewis DB, Trout KE. 2013. Hydrologic characterization of 56 geographically isolated wetlands in west-central Florida using a probabilistic method. *Wetland Ecology and Management* 21: 1–14.
- Ogawa H, Male JW. 1986. Simulating the flood mitigation role of wetlands. *Journal of Water Resources Planning and Management* 112: 114–128.
- Park J, Botter G, Jawitz JW, Rao PSC. 2014. Stochastic modeling of hydrologic variability of geographically isolated wetlands: effects of hydro-climatic forcing and wetland bathymetry. *Advances in Water Resources* 69: 38–48.
- Phillips RW, Spence C, Pomeroy JW. 2011. Connectivity and runoff dynamics in heterogeneous basins. *Hydrological Processes* 25: 3061–3075.
- Poff NL, Allan JD, Bain MB, Karr JR, Prestegard KL, Richter BD, Sparks RE, Stromberg JC. 1997. The natural flow regime. *BioScience* 47: 769–784.
- Pyke CR, Marty J. 2005. Cattle grazing mediates climate change impacts on ephemeral wetlands. *Conservation Biology* 19: 1619–1625.
- Pyzoza JE, Callahan TJ, Sun G, Trettin CC, Miwa M. 2008. A conceptual hydrologic model for a forested Carolina Bay depression wetland on the Coastal Plain of South Carolina, USA. *Hydrologic Processes* 22: 2689–2698.
- Rains MC. 2011. Water sources and hydrodynamics of closed-basin depressions, Cook Inlet Region, Alaska. *Wetlands* 31: 377–387.
- Rains MC, Fogg GE, Harter T, Dahlgren RA, Williamson RJ. 2006. The role of perched aquifers in hydrological connectivity and biogeochemical processes in vernal pool landscapes, Central Valley, California. *Hydrological Processes* 20: 1157–1175.
- Rains MC, Dahlgren RA, Williamson RJ, Fogg GE, Harter T. 2008. Geological control of physical and chemical hydrology in vernal pools, Central Valley, California. *Wetlands* 28: 347–362.
- Randall GW, Higgins DR, Russelle MP, Fuchs DJ, Nelson WW, Andrews JL. 1997. Nitrate losses through subsurface tile drainage in conservation reserve program, alfalfa, and row crop systems. *Journal of Environmental Quality* 26: 1240–1247.
- Rosenberry DO, Winter TC. 1997. Dynamics of water-table fluctuations in an upland between two prairie–pothole wetlands in North Dakota. *Journal of Hydrology* 191: 266–289.
- Sass GZ, Creed IF. 2008. Characterizing hydrodynamics on boreal landscapes using archived synthetic aperture radar imagery. *Hydrological Processes* 22: 1687–1699.
- Semlitsch RD, Bodie JR. 1998. Are small, isolated wetlands expendable? *Conservation Biology* 12: 1129–1133.
- Sinclair WC. 1977. Experimental study of artificial recharge possibilities in Northwest Hillsborough County, Florida, US Geological Survey Water-Resources Investigations 77–13.
- Sun RJ, Weeks JB, Grubb HF. 1997. Bibliography of regional aquifer-system analysis program of the US Geological Survey, 1978–96. US Geological Survey Water-Resources Investigations Report 97–4074.
- Sun G, McNulty SG, Amatya DM, Skaggs RW, Swift LW, Shepard JP, Riekerk H. 2002. A comparison of the hydrology of the coastal forested wetlands/pine flatwoods and the mountainous uplands in the southern US. *Journal of Hydrology* 263: 92–104.
- Tanner GW, Sandoval LD, Martin FG. 1984. Cattle behavior on a south Florida range. *Journal of Range Management* 37: 248–251.
- Tiner RW. 2003a. Estimated extent of geographically isolated wetlands in selected areas of the United States. *Wetlands* 23: 636–652.
- Tiner RW. 2003b. Geographically isolated wetlands of the United States. *Wetlands* 23: 494–516.
- Towler BW, Cahoon JE, Stein OR. 2004. Evapotranspiration crop coefficients for cattail and bulrush. *Journal of Hydrologic Engineering* 9: 235–239.
- Tsai J-S, Verne LS, McMurry ST, Smith LM. 2007. Influence of land use and wetland characteristics on water loss rates and hydroperiods of playas in the southern high plains. *Wetlands* 27: 683–692.
- US Environmental Protection Agency. 2015. *Connectivity of Streams and Wetlands to Downstream Waters: A Review and Synthesis of the Scientific Evidence*. EPA/600/R-14/475F, US Environmental Protection Agency Washington, DC.
- Van Meter KJ, Basu NB. 2015. Signatures of human impact: size distributions and spatial organization of wetlands in the prairie pothole landscape. *Ecological Applications* 25: 451–465.
- Walsh CJ, Fletcher TD, Burns MJ. 2012. Urban stormwater runoff: a new class of environmental flow problem. *PloS One* 7: e45814 doi:10.1371/journal.pone.0045814.

- Ward JV. 1989. The four-dimensional nature of lotic ecosystems. *Journal of the North American Benthological Society* 8: 2–8.
- Wigington PJ Jr, Leibowitz SG, Comeleo RL, Ebersole JL. 2013. Oregon hydrologic landscapes: a classification framework. *Journal of the American Water Resources Association* 49: 163–182.
- Wilcox B, Dean D, Jacob J, Sipocz A. 2011. Evidence of surface connectivity for Texas Gulf Coast depression wetlands. *Wetlands* 31: 451–458.
- Winter TC. 1999. Relation of streams, lakes, and wetlands to groundwater flow systems. *Hydrogeology Journal* 7: 28–45.
- Winter TC. 2001. The concept of hydrologic landscapes. *Journal of the American Water Resources Association* 37: 335–349.
- Wolock DM, Winter TC, McMahon G. 2004. Delineation and evaluation of hydrologic-landscape regions in the United States using geographic information system tools and multivariate statistical analyses. *Environmental Management* 34: S71–S88.
- Wood WW, Sanford WE. 1995. Chemical and isotopic methods for quantifying ground-water recharge in a regional, semiarid environment. *Ground Water* 33: 458–468.
- Yang J, Chu X. 2012. Effects of DEM resolution on surface depression properties and hydrologic connectivity. *Journal of Hydrologic Engineering* 18: 1157–1169.